

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau





(43) International Publication Date  
27 December 2002 (27.12.2002)

PCT

(10) International Publication Number  
**WO 02/102801 A1**

- (51) International Patent Classification<sup>7</sup>: **C07D 451/02**,  
A61K 31/46
- (74) Common Representative: **NEUROSEARCH A/S**,  
Patent Department, 93 Pederstrupvej, DK-2750 Ballerup  
(DK).
- (21) International Application Number: **PCT/DK02/00346**
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: **23 May 2002 (23.05.2002)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:  
PA 2001 00838 **23 May 2001 (23.05.2001)** **DK**
- (84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **NEU-  
ROSEARCH A/S [DK/DK]**; 93 Pederstrupvej, DK-2750  
Ballerup (DK).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **GOULIAEV, Alex**,  
**Haahr [DK/DK]**; Brøndsted 223, DK-3670 Veksø Sj.  
(DK). **DAHL, Bjarne, H. [DK/DK]**; NeuroSearch A/S, 93  
Pederstrupvej, DK-2750 Ballerup (DK). **PETERS, Dan**  
[SE/DK]; NeuroSearch A/S, 93 Pederstrupvej, DK-2750  
Ballerup (DK).
- Published:  
— *with international search report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **TROPANE DERIVATIVES AND THEIR USE AS MONOAMINE NEUROTRANSMITTER RE-UP TAKE INHIBITORS**



WO 02/102801 A1

(57) Abstract: This invention relates to tropane derivatives. In other aspects the invention relates to the use of these compounds in a method for therapy and to pharmaceutical compositions comprising the optically active isomer of the invention.

## TROPANE DERIVATIVES AND THEIR USE AS MONOAMINE NEUROTRANSMITTER RE-UP TAKE INHIBITORS

5

### TECHNICAL FIELD

This invention relates to tropane derivatives.

In other aspects the invention relates to the use of these compounds in a method for therapy and to pharmaceutical compositions comprising the optically  
10 active isomer of the invention.

### BACKGROUND ART

EP 604355, EP 604352, US 5444070, EP 604354, and WO 97/30997  
15 describe tropane derivatives and their use as mixed monoamine neurotransmitter re-uptake inhibitors. The documents specifically disclose a number of (1R,2R,3S)-2,3-disubstituted tropane derivatives and (1R,2S,3S)-2,3-disubstituted tropane derivatives.


However, there is a continued strong need to find compounds with an  
20 optimised biochemical profile as regards the activity on reuptake of the monoamine neurotransmitters serotonin, dopamine and noradrenaline, such as the ratio of the serotonin reuptake versus the noradrenaline and dopamine activity.

Furthermore, there is a strong need to find effective compounds, which structurally and synthetically wise are unrelated to cocaine.

25

### SUMMARY OF THE INVENTION

Therefore, in its first aspect, the invention provides a disubstituted tropane  
30 derivative of general formula Ia or Ib,



or a pharmaceutically acceptable addition salt thereof or the N-oxide thereof.


In its second aspect the invention provides a pharmaceutical composition comprising a therapeutically effective amount of a compound of the invention together with at least one pharmaceutically-acceptable carrier, excipient or diluent.

In a further aspect the invention provides a method of treatment, prevention  
5 or alleviation of a disease or a disorder or a condition of a living animal body, including a human, which disorder, disease or condition is responsive to inhibition of monoamine neurotransmitter re-uptake in the central nervous system (CNS), which method comprises the step of administering to such a living animal body in need thereof a therapeutically effective amount of a compound of the invention.

10 Other objects of the invention will be apparent to the person skilled in the art from the following detailed description and examples.

### DETAILED DISCLOSURE OF THE INVENTION

15 In its first aspect, the invention provides a disubstituted tropane derivative of general formula (Ia) or (Ib),



20

or a pharmaceutically acceptable addition salt thereof or the N-oxide thereof, wherein

R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl or 2-hydroxyethyl;

R<sup>3</sup> is

- 25     • **CH<sub>2</sub>-X-R'**,  
          wherein X is O, S, or NR'';  
          wherein R'' is hydrogen or alkyl; and  
          R' is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, or -CO-alkyl;
- 30     • **heteroaryl** which may be substituted one or more times with
- alkyl, cycloalkyl, or cycloalkylalkyl;
  - phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;

- phenylphenyl;
  - pyridyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
  - 5      ○ thienyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl; or
  - benzyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl; or
  - 10      • (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sup>11</sup>, COR<sup>11</sup>, or CH<sub>2</sub>R<sup>12</sup>;  
       wherein R<sup>11</sup> is
    - alkyl, cycloalkyl, or cycloalkylalkyl;
    - phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
    - 15      ○ phenylphenyl;
    - pyridyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
    - 20      ○ thienyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl; or
    - benzyl;
  - 25      n is 0 or 1; and  
       R<sup>12</sup> is
    - O-phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl; or
    - 30      ○ O-CO-phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
- R<sup>4</sup> is
  - 3,4-methylenedioxyphenyl or
  - 35      • phenyl, benzyl, naphthyl, or heteroaryl all of which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, cycloalkoxy, alkyl, cycloalkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl.

In its second aspect, the invention provides a pharmaceutical composition, comprising a therapeutically effective amount of a compound of the invention, or a pharmaceutically acceptable addition salt thereof, together with at least one pharmaceutically acceptable carrier, excipient or diluent.

5 In a further aspect, the invention provides the use of a compound of the invention, or a pharmaceutically acceptable addition salt thereof, for the manufacture of a pharmaceutical composition for the treatment, prevention or alleviation of a disease or a disorder or a condition of a mammal, including a human, which disease, disorder or condition is responsive to inhibition of monoamine neurotransmitter re-  
10 uptake in the central nervous system.

In a still further aspect, the invention relates to a method for treatment, prevention or alleviation of a disease or a disorder or a condition of a living animal body, including a human, which disorder, disease or condition is responsive to inhibition of monoamine neurotransmitter re-uptake in the central nervous system,  
15 which method comprises the step of administering to such a living animal body in need thereof a therapeutically effective amount of a compound of the invention.

In one embodiment, the invention relates to compounds of formula (Ia). In a second embodiment, the invention relates to compounds of formula (Ib). In a further embodiment, the invention relates to (1S,2S,3R,5R)-8-aza-bicyclo[3.2.1]octane  
20 derivatives. In a still further embodiment, the invention relates to (1S,2R,3R,5R)-8-aza-bicyclo[3.2.1]octane derivatives.

In a further embodiment, R<sup>3</sup> is

- **1,2,4-oxadiazol-3-yl** which may be substituted in the 5 position with
  - alkyl, cycloalkyl, or cycloalkylalkyl;
  - 25 ○ phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
  - phenylphenyl; or
  - benzyl which may be substituted one or more times with substituents  
30 selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl; or
- **1,2,4-oxadiazol-5-yl** which may be substituted in the 3 position with
  - alkyl, cycloalkyl, or cycloalkylalkyl;
  - 35 ○ phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
  - phenylphenyl;

- benzyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
- pyridyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro and heteroaryl; or
- thienyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro and heteroaryl.

In still further embodiment, R<sup>3</sup> is

- **CH<sub>2</sub>-X-R'**,  
wherein X is O, S, or NR'';  
wherein R'' is hydrogen or alkyl; and  
R' is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, or -CO-alkyl.

In a further embodiment, R<sup>4</sup> is

- **phenyl**, which is substituted once or twice with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, cycloalkoxy, alkyl, cycloalkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl.

In a further embodiment, R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, or 2-hydroxyethyl;

R<sup>3</sup> is

- **CH<sub>2</sub>-X-R'**,  
wherein X is O, S, or NR'';  
wherein R'' is hydrogen or alkyl; and  
R' is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, or -CO-alkyl; or
- **(CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sup>11</sup>, or COR<sup>11</sup>;**  
wherein R<sup>11</sup> is alkyl, cycloalkyl, or cycloalkylalkyl; and n is 0 or 1;

R<sup>4</sup> is

- **phenyl** which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, cycloalkoxy, alkyl, cycloalkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl.

In a still further embodiment, R is hydrogen or alkyl;

R<sup>3</sup> is

- **CH<sub>2</sub>-X-R'**,  
wherein X is O or S; and  
R' is hydrogen or alkyl; or
- **(CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sup>11</sup>;**  
wherein R<sup>11</sup> is alkyl; and n is 0 or 1;

R<sup>4</sup> is

- **phenyl** which may be substituted once or twice with substituents selected from the group consisting of halogen, CF<sub>3</sub>, and CN.

In a special embodiment, R is hydrogen. In a further embodiment, R is alkyl, such as methyl.

- 5 In a further special embodiment, R<sup>3</sup> is CH<sub>2</sub>-X-R', wherein X is O; and R' is hydrogen or alkyl. In one embodiment, R' is hydrogen. In a second embodiment, R' is methyl. In a further embodiment, R' is ethyl.

- In a special embodiment, R<sup>4</sup> is phenyl which may be substituted once or twice with substituents selected from the group consisting of halogen, CF<sub>3</sub>, and CN. In one  
10 embodiment R<sup>4</sup> is phenyl which may be substituted once or twice with halogen, such as chlorine. In a second embodiment, R<sup>4</sup> is phenyl substituted once or twice with chlorine. In a special embodiment, R<sup>4</sup> is phenyl 3,4-dichlorophenyl.

In a special embodiment, the compound of general formula I is selected from:

- 15 (1S,2R,3R,5R)-3-(3,4-Dichlorophenyl)-8-methyl-8-aza-bicyclo[3.2.1]octane-2-carboxylic acid ethyl ester;  
(1S,2S,3R,5R)-3-(3,4-Dichlorophenyl)-8-methyl-8-aza-bicyclo[3.2.1]octane-2-carboxylic acid ethyl ester;  
(1S,2R,3R,5R)-2-Hydroxymethyl-3-(3,4-dichlorophenyl)-tropane;  
20 (1S,2S,3R,5R)-2-Hydroxymethyl-3-(3,4-dichlorophenyl)-tropane;  
(1S,2R,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-tropane;  
(1S,2S,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-tropane;  
(1S,2R,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-8H-8-aza-bicyclo[3.2.1]octane;  
(1S,2S,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-8H-8-aza-bicyclo[3.2.1]octane;  
25 or a pharmaceutically acceptable addition salt thereof.

### Definition of Substituents

In the context of this invention halogen represents a fluorine, a chlorine, a bromine or an iodine atom.

- 30 Alkyl means a straight chain or branched chain of one to six carbon atoms, including but not limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, pentyl, and hexyl; methyl, ethyl, propyl and isopropyl are preferred groups.

Cycloalkyl means cyclic alkyl of three to seven carbon atoms, including but not limited to cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl;

- 35 Alkenyl means a group of from two to six carbon atoms, including at least one double bond, for example, but not limited to ethenyl, 1,2- or 2,3-propenyl, or 1,2-, 2,3-, or 3,4-butenyl.

Alkynyl means a group of from two to six carbon atoms, including at least one triple bond, for example, but not limited to ethynyl, 1,2-, 2,3-propynyl, or 1,2-, 2,3- or 3,4-butynyl.

Alkoxy is O-alkyl, wherein alkyl is as defined above.

5 Acyl is -CO-alkyl wherein alkyl is as defined above.

Cycloalkoxy means O-cycloalkyl, wherein cycloalkyl is as defined above.

Cycloalkylalkyl means cycloalkyl as above and alkyl as above, meaning for example, cyclopropylmethyl.

Amino is  $\text{NH}_2$  or NH-alkyl or N-(alkyl)<sub>2</sub>, wherein alkyl is as defined above.

10 Aryl is a carbocyclic aromatic ring system such as phenyl or naphthyl (1-naphthyl or 2-naphthyl).

Heteroaryl is a 5- or 6-membered heterocyclic monocyclic group, for example, but not limited to, oxazol-2-yl, oxazol-4-yl, oxazol-5-yl, isoxazol-3-yl, isoxazol-4-yl, isoxazol-5-yl, thiazol-2-yl, thiazol-4-yl, thiazol-5-yl, isothiazol-3-yl, isothiazol-4-yl, 15 isothiazol-5-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl, 1,2,5-oxadiazol-3-yl, 1,2,5-oxadiazol-4-yl, 1,2,5-thiadiazol-3-yl, 1,2,5-thiadiazol-4-yl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 2-pyrrolyl, 3-pyrrolyl, 2-furanyl, 3-furanyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl.

20 The compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention.

The compounds of the invention may be prepared in numerous ways. The 25 compounds of the invention and their pharmaceutically acceptable derivatives may thus be prepared by any method known in the art for the preparation of compounds of analogous structure, and as shown in the representative examples which follow.

#### Pharmaceutically Acceptable Salts

30 The chemical compound of the invention may be provided in any form suitable for the intended administration. Suitable forms include pharmaceutically (i.e. physiologically) acceptable salts, and pre- or prodrug forms of the chemical compound of the invention.

Examples of pharmaceutically acceptable addition salts include, without 35 limitation, the non-toxic inorganic and organic acid addition salts such as the hydrochloride derived from hydrochloric acid, the hydrobromide derived from hydrobromic acid, the nitrate derived from nitric acid, the perchlorate derived from perchloric acid, the phosphate derived from phosphoric acid, the sulphate derived from sulphuric acid, the formate derived from formic acid, the acetate derived from



acetic acid, the aconate derived from aconitic acid, the ascorbate derived from ascorbic acid, the benzenesulphonate derived from benzenesulphonic acid, the benzoate derived from benzoic acid, the cinnamate derived from cinnamic acid, the citrate derived from citric acid, the embonate derived from embonic acid, the enantate  
5 derived from enanthic acid, the fumarate derived from fumaric acid, the glutamate derived from glutamic acid, the glycolate derived from glycolic acid, the lactate derived from lactic acid, the maleate derived from maleic acid, the malonate derived from malonic acid, the mandelate derived from mandelic acid, the methanesulphonate derived from methane sulphonic acid, the naphthalene-2-sulphonate derived from  
10 naphthalene-2-sulphonic acid, the phthalate derived from phthalic acid, the salicylate derived from salicylic acid, the sorbate derived from sorbic acid, the stearate derived from stearic acid, the succinate derived from succinic acid, the tartrate derived from tartaric acid, the toluene-p-sulphonate derived from p-toluene sulphonic acid, and the like. Such salts may be formed by procedures well known and described in the art.

15 Other acids such as oxalic acid, which may not be considered pharmaceutically acceptable, may be useful in the preparation of salts useful as intermediates in obtaining a chemical compound of the invention and its pharmaceutically acceptable acid addition salt.

Metal salts of a chemical compound of the invention includes alkali metal  
20 salts, such as the sodium salt of a chemical compound of the invention containing a carboxy group.

In the context of this invention the "onium salts" of N-containing compounds are also contemplated as pharmaceutically acceptable salts. Preferred "onium salts" include the alkyl-onium salts, the cycloalkyl-onium salts, and the  
25 cycloalkylalkyl-onium salts.

The chemical substance according to the invention may be administered as such or in the form of a suitable prodrug.

The term "prodrug" denotes a compound, which is a drug precursor and which, following administration and absorption, release the drug in vivo via some  
30 metabolic process.

Particularly favoured prodrugs are those that increase the bioavailability of the compounds of the invention (e.g. by allowing an orally administered compound to be more readily absorbed into the blood) or which enhance delivery of the parent compound to a specific biological compartment (e.g. the brain or lymphatic system).

35 Thus examples of suitable prodrugs of the substances according to the invention include compounds modified at one or more reactive or derivatizable groups of the parent compound. Of particular interest are compounds modified at a carboxyl group, a hydroxyl group, or an amino group. Examples of suitable derivatives are esters or amides.

### Labelled Compounds

The compounds of the invention may be used in their labelled or unlabelled form. In the context of this invention "label" stands for the binding of a marker to the  
5 compound of interest that will allow easy quantitative detection of said compound.

The labelled compounds of the invention may be useful as diagnostic tools, radio tracers, or monitoring agents in various diagnostic methods, and for *in vivo* receptor imaging.

The labelled isomer of the invention preferably contains at least one  
10 radionuclide as a label. Positron emitting radionuclides are all candidates for usage. In the context of this invention the radionuclide is preferably selected from  $^2\text{H}$  (deuterium),  $^3\text{H}$  (tritium),  $^{13}\text{C}$ ,  $^{14}\text{C}$ ,  $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^{123}\text{I}$ , and  $^{18}\text{F}$ .

The physical method for detecting the labelled isomer of the present invention may be selected from Position Emission Tomography (PET), Single Photon  
15 Imaging Computed Tomography (SPECT), Magnetic Resonance Spectroscopy (MRS), Magnetic Resonance Imaging (MRI), and Computed Axial X-ray Tomography (CAT), or combinations thereof.

### **Methods of Preparation**

20 The compounds of the invention may be prepared by conventional methods for chemical synthesis, e.g. those described in the working examples. The starting materials for the processes described in the present application are known or may readily be prepared by conventional methods from commercially available chemicals.

25 Also one compound of the invention can be converted to another compound of the invention using conventional methods, such as those described in EP 604355, EP 604352, US 5444070, EP 604354, and WO 97/30997.

The end product of the reaction described herein may be isolated by conventional techniques, e.g. by extraction, crystallisation, distillation,  
30 chromatography, etc.

### **Biological Activity**

The compounds of the invention may be tested for its ability to inhibit the reuptake of the monoamine neurotransmitters in synaptosomes, eg such as  
35 described in WO 97/30997. Based on the balanced activity observed in these tests the compound of the invention is considered useful for combating diseases, disorders or conditions associated with the dopaminergic, noradrenalinergic and/or serotonergic neural system.

The diseases, disorders or conditions contemplated in this context are eating disorders, obesity, anorexia nervosa, disorders of sleep, panic disorders, social phobia, dementia, senile dementia, pre-senile dementia, memory deficits, memory loss, Alzheimer's disease, chronic fatigue syndrome, anxiety, pseudodementia, Ganser's syndrome, narcolepsy, drug addiction or misuse including cocaine abuse, alcoholism, tobacco abuse, panic disorder, post-traumatic syndrome, migraine, pain, attention deficit hyperactivity disorder, autism, mutism, trichotillomania, Parkinson's disease, depression, attention, alertness, arousal, vigilance, premature ejaculation, and erectile dysfunction.

10 The compounds of the invention are considered particularly useful for the treatment, prevention or alleviation of depression, pseudodementia, Ganser's syndrome, obsessive compulsive disorders, panic disorders, memory deficits, attention deficit hyperactivity disorder, obesity, anxiety, and eating disorders.

#### 15 **Pharmaceutical Compositions**

In another aspect the invention provides novel pharmaceutical compositions comprising a therapeutically effective amount of the chemical compound of the invention.

20 While a chemical compound of the invention for use in therapy may be administered in the form of the raw chemical compound, it is preferred to introduce the active ingredient, optionally in the form of a physiologically acceptable salt, in a pharmaceutical composition together with one or more adjuvants, excipients, carriers, buffers, diluents, and/or other customary pharmaceutical auxiliaries.

In a preferred embodiment, the invention provides pharmaceutical compositions comprising the chemical compound of the invention, or a pharmaceutically acceptable salt or derivative thereof, together with one or more pharmaceutically acceptable carriers therefor, and, optionally, other therapeutic and/or prophylactic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not harmful to the recipient thereof.

30 Pharmaceutical compositions of the invention may be those suitable for oral, rectal, bronchial, nasal, topical (including buccal and sub-lingual), transdermal, vaginal or parenteral (including cutaneous, subcutaneous, intramuscular, intraperitoneal, intravenous, intraarterial, intracerebral, intraocular injection or infusion) administration, or those in a form suitable for administration by inhalation or insufflation, including powders and liquid aerosol administration, or by sustained release systems. Suitable examples of sustained release systems include semipermeable matrices of solid hydrophobic polymers containing the compound of

the invention, which matrices may be in form of shaped articles, e.g. films or microcapsules.

The chemical compound of the invention, together with a conventional adjuvant, carrier, or diluent, may thus be placed into the form of pharmaceutical compositions and unit dosages thereof. Such forms include solids, and in particular tablets, filled capsules, powder and pellet forms, and liquids, in particular aqueous or non-aqueous solutions, suspensions, emulsions, elixirs, and capsules filled with the same, all for oral use, suppositories for rectal administration, and sterile injectable solutions for parenteral use. Such pharmaceutical compositions and unit dosage forms thereof may comprise conventional ingredients in conventional proportions, with or without additional active compounds or principles, and such unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed.

The chemical compound of the present invention can be administered in a wide variety of oral and parenteral dosage forms. It will be obvious to those skilled in the art that the following dosage forms may comprise, as the active component, either a chemical compound of the invention or a pharmaceutically acceptable salt of a chemical compound of the invention.

For preparing pharmaceutical compositions from a chemical compound of the present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavouring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

In powders, the carrier is a finely divided solid, which is in a mixture with the finely divided active component.

In tablets, the active component is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired.

The powders and tablets preferably contain from five or ten to about seventy percent of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as carrier providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders,

capsules, pills, cachets, and lozenges can be used as solid forms suitable for oral administration.

For preparing suppositories, a low melting wax, such as a mixture of fatty acid glyceride or cocoa butter, is first melted and the active component is dispersed  
5 homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized moulds, allowed to cool, and thereby to solidify.

Compositions suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

10 Liquid preparations include solutions, suspensions, and emulsions, for example, water or water-propylene glycol solutions. For example, parenteral injection liquid preparations can be formulated as solutions in aqueous polyethylene glycol solution.

The chemical compound according to the present invention may thus be  
15 formulated for parenteral administration (e.g. by injection, for example bolus injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain formulation agents such as  
20 suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilization from solution, for constitution with a suitable vehicle, e.g. sterile, pyrogen-free water, before use.

Aqueous solutions suitable for oral use can be prepared by dissolving the  
25 active component in water and adding suitable colorants, flavours, stabilising and thickening agents, as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, or other well  
30 known suspending agents.

Also included are solid form preparations, intended for conversion shortly before use to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. In addition to the active component such preparations may comprise colorants, flavours, stabilisers, buffers, artificial and  
35 natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

For topical administration to the epidermis the chemical compound of the invention may be formulated as ointments, creams or lotions, or as a transdermal patch. Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be

formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilising agents, dispersing agents, suspending agents, thickening agents, or colouring agents.

Compositions suitable for topical administration in the mouth include  
5 lozenges comprising the active agent in a flavoured base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert base such as gelatin and glycerine or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Solutions or suspensions are applied directly to the nasal cavity by  
10 conventional means, for example with a dropper, pipette or spray. The compositions may be provided in single or multi-dose form.

Administration to the respiratory tract may also be achieved by means of an aerosol formulation in which the active ingredient is provided in a pressurised pack with a suitable propellant such as a chlorofluorocarbon (CFC) for example  
15 dichlorodifluoromethane, trichlorofluoromethane, or dichlorotetrafluoroethane, carbon dioxide, or other suitable gas. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by provision of a metered valve.

Alternatively the active ingredients may be provided in the form of a dry powder, for example a powder mix of the compound in a suitable powder base such  
20 as lactose, starch, starch derivatives such as hydroxypropylmethyl cellulose and polyvinylpyrrolidone (PVP). Conveniently the powder carrier will form a gel in the nasal cavity. The powder composition may be presented in unit dose form for example in capsules or cartridges of, e.g., gelatin, or blister packs from which the powder may be administered by means of an inhaler.

25 In compositions intended for administration to the respiratory tract, including intranasal compositions, the compound will generally have a small particle size for example of the order of 5 microns or less. Such a particle size may be obtained by means known in the art, for example by micronization.

When desired, compositions adapted to give sustained release of the  
30 active ingredient may be employed.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as  
35 packaged tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

Tablets or capsules for oral administration and liquids for intravenous administration and continuous infusion are preferred compositions.

Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

A therapeutically effective dose refers to that amount of active ingredient, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity, e.g. ED<sub>50</sub> and LD<sub>50</sub>, may be determined by standard pharmacological procedures in cell cultures or experimental animals. The dose ratio between therapeutic and toxic effects is the therapeutic index and may be expressed by the ratio LD<sub>50</sub>/ED<sub>50</sub>. Pharmaceutical compositions exhibiting large therapeutic indexes are preferred.

10 The dose administered must of course be carefully adjusted to the age, weight and condition of the individual being treated, as well as the route of administration, dosage form and regimen, and the result desired, and the exact dosage should of course be determined by the practitioner.

The actual dosage depend on the nature and severity of the disease being treated and the route of administration, and is within the discretion of the physician, and may be varied by titration of the dosage to the particular circumstances of this invention to produce the desired therapeutic effect. However, it is presently contemplated that pharmaceutical compositions containing of from about 0.01 to about 500 mg of active ingredient per individual dose, preferably of from about 0.1 to about 100 mg, most preferred of from about 1 to about 10 mg, are suitable for therapeutic treatments.

The active ingredient may be administered in one or several doses per day. A satisfactory result can, in certain instances, be obtained at a dosage as low as 0.01 µg/kg i.v. and 0.1 µg/kg p.o. The upper limit of the dosage range is presently considered to be about 10 mg/kg i.v. and 100 mg/kg p.o. Preferred ranges are from about 0.1 µg/kg to about 10 mg/kg/day i.v., and from about 1 µg/kg to about 100 mg/kg/day p.o.

### Methods of Therapy

30 In another aspect the invention provides a method for the treatment, prevention or alleviation of a disease or a disorder or a condition of a living animal body, including a human, which disease, disorder or condition is responsive to inhibition of monoamine neurotransmitter re-uptake in the central nervous system (CNS), and which method comprises administering to such a living animal body, including a human, in need thereof an effective amount of the optically active isomer of the invention.

In a more preferred embodiment the invention provides a method of combating depression, pseudodementia, Ganser's syndrome, obsessive compulsive

disorders, panic disorders, memory deficits, attention deficit hyperactivity disorder, obesity, anxiety and eating disorders.


It is at present contemplated that suitable dosage ranges are 0.1 to 1000 milligrams daily, 10-500 milligrams daily, and especially 30-100 milligrams daily, dependent as usual upon the exact mode of administration, form in which administered, the indication toward which the administration is directed, the subject involved and the body weight of the subject involved, and further the preference and experience of the physician or veterinarian in charge. A satisfactory result can, in certain instances, be obtained at a dosage as low as 0.005 mg/kg i.v. and 0.01 mg/kg p.o. The upper limit of the dosage range is about 10 mg/kg i.v. and 100 mg/kg p.o. Preferred ranges are from about 0.001 to about 1 mg/kg i.v. and from about 0.1 to about 10 mg/kg p.o.

Any possible combination of two or more of the embodiments described herein is comprised within the scope of the present invention.

## EXAMPLES

The invention is further illustrated with reference to the following examples, which are not intended to be in any way limiting to the scope of the invention as claimed.


### Example 1



1

**(-)-2-Carbomethoxytropinone (1):** Was prepared by a known procedure (J. F. Casale, Forensic Science International, 33 (1987) 275-298).

### Example 2





2



**(+)-Ecgonine ethylester (2):** To a stirred solution of (-)-2-carbomethoxytropinone (37.4 g, 0.19 mol) in methanol (1.5 l) at  $-45^{\circ}\text{C}$ , was added sodium borohydride (37.0 g, 0.98 mol) in small portions, such that the internal temperature was kept between  $-45^{\circ}\text{C}$  and  $-35^{\circ}\text{C}$ . The reaction mixture was stirred at  $-45^{\circ}\text{C}$  for 2 hours, and  
5 quenched by dropwise addition of concentrated hydrochloric acid (120 ml), while keeping the temperature at  $-45^{\circ}\text{C}$ . The reaction mixture was allowed to warm to room temperature and stirred overnight. The reaction mixture was concentrated to a volume of approximately 120 ml, water (500 ml) was added and washed with diethyl ether (3 x 100 ml). To the aqueous phase was added aqueous ammonia (25%), until  
10 basic reaction, and extracted with dichloromethane (4 x 200 ml). The combined organic phases were dried with sodium sulphate and evaporated to an oil. The oil was dissolved in ethyl acetate (370 ml) and a solution of sodium ethoxide in ethanol (300 ml, 1 M, prepared from sodium (7.0 g, 303 mmol), was added. The resulting solution was heated at reflux for 3 hours, cooled to room-temperature and evaporated to an  
15 oil. The residue was solved in toluene (0.5 l) and evaporated to an oil, this was repeated. The product 30 g (79%) was obtained as an oil.


### Example 3



**(1S,2R,3R,5R)-3-(3,4-Dichlorophenyl)-8-methyl-8-aza-bicyclo[3.2.1]octane-2-carboxylic acid ethyl ester (3) and (1S,2S,3R,5R)-3-(3,4-dichlorophenyl)-8-methyl-8-aza-bicyclo[3.2.1]octane-2-carboxylic acid ethyl ester (4):** A stirred  
25 suspension of magnesium turnings (1.25 g, 52 mmol) in diethyl ether was added bromo-3,4-dichlorobenzene (11.5 g, 49 mmol). The mixture was heated at reflux for 30 minutes and then cooled to  $-20^{\circ}\text{C}$ . A solution of (+)-ecgonine ethylester (5.0 g, 25.6 mmol) in toluene (30 ml) was slowly added, while keeping the internal temperature between  $-15^{\circ}\text{C}$  and  $-10^{\circ}\text{C}$ . The reaction mixture was stirred at  $-15^{\circ}\text{C}$  for  
30  $1\frac{1}{2}$  hour and then added trifluoroacetic acid (8.0 ml) and water (100 ml). Concentrated hydrochloric acid was added until  $\text{pH} = 1$ . The phases were separated and the aqueous phase was washed with diethyl ether (2 x 100 ml). The aqueous phase was added 25% aqueous ammonia until  $\text{pH} = 11$  and extracted with dichloromethane (2 x 100 ml). The combined organic phases were dried using magnesium sulphate and  
35 then evaporated to an oil. Yield 7.5 g (86 %) of (3) and (4). The isomers (3) and (4)

were separated using column chromatography and petroleum ether, diethyl ether, triethylamine (1 : 1 : 2 %) to give 3.5 g (40 %) of (3), Mp 67.5-68.5°C and 1.7 g (20 %) of (4).


#### 5 Example 4



##### Method A

**(1S,2R,3R,5R)-2-Hydroxymethyl-3-(3,4-dichlorophenyl)-tropane (5):** A stirred solution of (1S,2R,3R,5R)-3-(3,4-dichlorophenyl)-8-methyl-8-aza-bicyclo[3.2.1]octane-2-carboxylic acid ethyl ester (3), (6.8 g, 20 mmol) in tetrahydrofuran (100 ml) at -40°C was added lithium aluminum hydride (LiAlH<sub>4</sub>) (1.0 g, 26 mmol), while keeping the internal temperature between -40-(-30) °C. The reaction mixture was left with stirring at -40 °C for 1 h, then quenched by addition of water (10 ml) followed by aqueous sodium hydroxide (10 ml, 4 M). The mixture was filtered and evaporated to an oil. The oil was dissolved in dichloromethane and dried with magnesium sulphate and evaporated to a solid. Yield 5.3 g (88 %), Mp. 81-86°C .

#### Example 5




**(1S,2S,3R,5R)-2-Hydroxymethyl-3-(3,4-dichlorophenyl)-tropane (6):** (1S,2S,3R,5R)-3-(3,4-dichlorophenyl)-8-methyl-8-aza-bicyclo[3.2.1]octane-2-carboxylic acid ethyl ester (4) ( 5.6 g, 17.1 mmol) was reduced according to method A giving 4.4 g (86 %) of the title compound.

18

**Example 6**

5


**Method B**

**(1S,2R,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-tropane (7):** (1S,2R,3R,5R)-2-Hydroxymethyl-3-(3,4-dichlorophenyl)-tropane (5) (1.0 g) in dry tetrahydrofuran (15 ml) was added sodium hydride (0.25g, 6.25 mmol) and stirred for 15 minutes. Diethyl sulfate (0.54 ml, 4.1 mmol) was added and the mixture stirred at 50°C for 3 h. The reaction mixture was poured into water and extracted twice, using diethyl ether (2 x 50 ml). The organic phases were dried by magnesium sulphate and evaporated to dryness. Column chromatography, using a mixture of dichloromethane, methanol and ammonia (9 : 1 : 1 %) yielded 200 mg (18 %) of the title compound as the free base. Mp. 52.5-54.5°C.

20

**Example 7**

25




30

**(1S,2S,3R, 5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-tropane citrate (8):** Was synthesized from (1S,2S,3R,5R)-2-hydroxymethyl-3-(3,4-dichlorophenyl)-tropane (6) (1.5 g) according to method B. Yield 290 mg as the free base. The free base was dissolved in ethanol (10 ml, 96 %) and added citric acid (190 mg, 1.0 mmol). The suspension was heated to obtain a clear solution and left for precipitation. The precipitate was isolated by filtration to yield 290 mg (11 %) of the title compound. Mp. 161.9 – 162.3°C

19

**Example 8**

5




9

**10 (1S,2R,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-8H-8-azabicyclo[3.2.1]octane fumarate (9):**

To a mixture of (1S,2R,3R,5R)-2-ethoxymethyl-3-(3,4-dichlorophenyl)-tropane (7) (0.5 g, 1.53 mmol) and toluene (5 ml), was added 1-chloroethyl chloroformate (0.25 ml, 2.3 mmol). The mixture was stirred at 100 °C for 48 hours. Water (30 ml) was added and the reaction mixture stirred for 6 hours at 75 °C. The mixture was extracted with diethyl ether (2 x 50 ml). The organic phase was dried with magnesium sulphate and evaporated. Column chromatography, using dichloromethane, methanol and ammonia (9, 1, 1 %). Yield 0.33 g (69 %). A solution of fumaric acid (4.1 ml, 0.078 M) in diethyl ether and MeOH (9 : 1) was added to the free base (100 mg, 0.32 mmol). The mixture was stirred for one hour. The precipitate was isolated by filtration yielding 110 mg (0.25 mmol) of the title compound. Mp 145-147.5 °C.

**25 Example 9**

30



10

**(1S,2S,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-8H-8-azabicyclo[3.2.1]octane fumarate (10):** The compound was synthesized from (1S,2S,3R,5R)-2-ethoxymethyl-3-(3,4-dichlorophenyl)-tropane (8) according to method B. Mp 219-220 °C.

**Example 10**

The following compounds and pharmaceutically acceptable salts thereof is prepared analogously using the methods as described above and in EP 604355, EP 604352, US 5444070, EP 604354, and WO 97/30997.

5

(1S,2S,3R,5R)-2-(3-Cyclopropyl-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)tropane;

(1S,2S,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)tropane;

(1S,2S,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(4-methylphenyl)-tropane;

(1S,2S,3R,5R)-2-(3-Benzyl-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)tropane;

10 

(1S,2S,3R,5R)-2-(3-(4-Phenyl-phenyl)-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)tropane;

(1S,2S,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(2-naphthyl)tropane;

(1S,2R,3R,5R)-2-(3-Cyclopropyl-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)tropane;

(1S,2R,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)tropane;

(1S,2R,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(4-methylphenyl)-tropane;

15 

(1S,2R,3R,5R)-2-(3-Benzyl-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)tropane;

(1S,2R,3R,5R)-2-(3-(4-Phenyl-phenyl)-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)tropane;

(1S,2R,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(2-naphthyl)tropane;

(1S,2S,3R,5R)-2-Methoxymethyl-3-(3,4-dichlorophenyl)-tropane;

(1S,2S,3R,5R)-2-Isopropoxymethyl-3-(3,4-dichlorophenyl)-tropane;

20 

(1S,2S,3R,5R)-2-Cyclopropylmethyloxymethyl-3-(3,4-dichlorophenyl)-tropane;

(1S,2S,3R,5R)-2-Methoxymethyl-3-(4-chlorophenyl)-tropane;

(1S,2S,3R,5R)-N-Normethyl-2-methoxymethyl-3-(4-chlorophenyl)-tropane;

(1S,2S,3R,5R)-2-Ethoxymethyl-3-(4-chlorophenyl)-tropane;

(1S,2S,3R,5R)-N-Normethyl-2-methoxymethyl-3-(3,4-dichlorophenyl)-tropane;

25 

(1S,2S,3R,5R)-N-Normethyl-2-ethoxymethyl-3-(4-chlorophenyl)-tropane;

(1S,2S,3R,5R)-N-Normethyl-2-cyclopropylmethyloxymethyl-3-(4-chlorophenyl)-tropane;

(1S,2S,3R,5R)-2-Cyclopropylmethyloxymethyl-3-(4-chlorophenyl)-tropane;

(1S,2S,3R,5R)-2-Ethylthiomethyl-3-(3,4-dichlorophenyl)-tropane;

(1S,2S,3R,5R)-2-Hydroxymethyl-3-(4-fluorophenyl)tropane;

30 

(1S,2S,3R,5R)-2-Hydroxymethyl-3-(3,4-dichlorophenyl)tropane;

(1S,2S,3R,5R)-N-Normethyl-N-(*tert*-butoxycarbonyl)-2-hydroxymethyl-3-(3,4-dichlorophenyl)tropane;

(1S,2S,3R,5R)-2-Hydroxymethyl-3-(4-chlorophenyl)tropane;

(1S,2R,3R,5R)-2-Methoxymethyl-3-(3,4-dichlorophenyl)-tropane;

35 

(1S,2R,3R,5R)-2-Isopropoxymethyl-3-(3,4-dichlorophenyl)-tropane;

(1S,2R,3R,5R)-2-Cyclopropylmethyloxymethyl-3-(3,4-dichlorophenyl)-tropane;

(1S,2R,3R,5R)-2-Methoxymethyl-3-(4-chlorophenyl)-tropane;

(1S,2R,3R,5R)-N-Normethyl-2-methoxymethyl-3-(4-chlorophenyl)-tropane;

(1S,2R,3R,5R)-2-Ethoxymethyl-3-(4-chlorophenyl)-tropane;

- (1S,2R,3R,5R)-N-Normethyl-2-methoxymethyl-3-(3,4-dichlorophenyl)-tropane;  
(1S,2R,3R,5R)-N-Normethyl-2-ethoxymethyl-3-(4-chlorophenyl)-tropane;  
(1S,2R,3R,5R)-N-Normethyl-2-cyclopropylmethyloxymethyl-3-(4-chlorophenyl)-tropane;  
(1S,2R,3R,5R)-2-Cyclopropylmethyloxymethyl-3-(4-chlorophenyl)-tropane;  
5 (1S,2R,3R,5R)-2-Ethylthiomethyl-3-(3,4-dichlorophenyl)-tropane;  
(1S,2R,3R,5R)-2-Hydroxymethyl-3-(4-fluorophenyl)tropane;  
(1S,2R,3R,5R)-2-Hydroxymethyl-3-(3,4-dichlorophenyl)tropane;  
(1S,2R,3R,5R)-N-Normethyl-N-(*tert*-butoxycarbonyl)-2-hydroxymethyl-3-(3,4-dichlorophenyl)tropane;  
10 (1S,2R,3R,5R)-2-Hydroxymethyl-3-(4-chlorophenyl)tropane;  
(1S,2S,3R,5R)-2-(3-(2-Furanyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2S,3R,5R)-2-(3-(3-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2S,3R,5R)-N-Normethyl-N-allyl-2-(3-(4-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
15 (1S,2S,3R,5R)-N-Normethyl-N-ethyl-2-(3-(4-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2S,3R,5R)-N-Normethyl-N-(2-hydroxyethyl) -2-(3-(4-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2S,3R,5R)-N-Normethyl-2-(3-(4-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
20 (1S,2S,3R,5R)-N-Normethyl-N-allyl-2-(3-(3-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2S,3R,5R)-N-Normethyl-N-allyl-2-(3-(2-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
25 (1S,2S,3R,5R)-2-(3-(2-Thienyl)-1,2,4-oxadiazol-5-yl)-3-(4-chlorophenyl)-tropane;  
(1S,2S,3R,5R)-2-(3-(2-Thienyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2S,3R,5R)-2-(3-(4-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2S,3R,5R)-2-(3-(2-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2S,3R,5R)-2-(3-(4-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(4-chlorophenyl)-tropane;  
30 (1S,2S,3R,5R)-2-(3-(3-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(4-chlorophenyl)-tropane;  
(1S,2S,3R,5R)-2-(3-(2-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(4-chlorophenyl)-tropane;  
(1S,2R,3R,5R)-2-(3-(2-Furanyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2R,3R,5R)-2-(3-(3-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2R,3R,5R)-N-Normethyl-N-allyl-2-(3-(4-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
35 (1S,2R,3R,5R)-N-Normethyl-N-ethyl-2-(3-(4-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2R,3R,5R)-N-Normethyl-N-(2-hydroxyethyl) -2-(3-(4-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;


- (1S,2R,3R,5R)-N-Normethyl-2-(3-(4-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;
- (1S,2R,3R,5R)-N-Normethyl-N-allyl-2-(3-(3-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;
- 5 (1S,2R,3R,5R)-N-Normethyl-N-allyl-2-(3-(2-pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;
- (1S,2R,3R,5R)-2-(3-(2-Thienyl)-1,2,4-oxadiazol-5-yl)-3-(4-chlorophenyl)-tropane;
- (1S,2R,3R,5R)-2-(3-(2-Thienyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;
- (1S,2R,3R,5R)-2-(3-(4-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;
- 10 (1S,2R,3R,5R)-2-(3-(2-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(3,4-dichlorophenyl)-tropane;
- (1S,2R,3R,5R)-2-(3-(4-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(4-chlorophenyl)-tropane;
- (1S,2R,3R,5R)-2-(3-(3-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(4-chlorophenyl)-tropane;
- (1S,2R,3R,5R)-2-(3-(2-Pyridyl)-1,2,4-oxadiazol-5-yl)-3-(4-chlorophenyl)-tropane;
- (1S,2S,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)-tropane;
- 15 (1S,2S,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(4-methylphenyl)-tropane;
- (1S,2S,3R,5R)-2-(3-Benzyl-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)-tropane;
- (1S,2S,3R,5R)-2-(3-(4-Phenylphenyl)-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)-tropane;
- (1S,2S,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(2-naphthyl)-tropane;
- (1S,2R,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)-tropane;
- 20 (1S,2R,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(4-methylphenyl)-tropane;
- (1S,2R,3R,5R)-2-(3-Benzyl-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)-tropane;
- (1S,2R,3R,5R)-2-(3-(4-Phenylphenyl)-1,2,4-oxadiazol-5-yl)-3-(4-fluorophenyl)-tropane;
- (1S,2R,3R,5R)-2-(3-Phenyl-1,2,4-oxadiazol-5-yl)-3-(2-naphthyl)-tropane;
- (1S,2S,3R,5R)-2-(4-Chlorophenoxy-methyl)-3-(4-fluorophenyl)-tropane;
- 25 (1S,2S,3R,5R)-2-(4-Chlorophenoxy-methyl)-3-(4-fluorophenyl)-tropane;
- (1S,2S,3R,5R)-2-(4-Chlorophenoxy-methyl)-3-(3,4-dichlorophenyl)-tropane;
- (1S,2S,3R,5R)-2-(4-Chlorophenoxy-methyl)-3-(4-methylphenyl)-tropane;
- (1S,2S,3R,5R)-2-(4-Benzoyloxy-methyl)-3-(4-fluorophenyl)-tropane;
- (1S,2S,3R,5R)-2-Carbomethoxy-3-(2-naphthyl)-tropane;
- 30 (1S,2S,3R,5R)-2-Carbomethoxy-3-(3,4-dichlorophenyl)-tropane;
- (1S,2S,3R,5R)-2-Carbomethoxy-3-benzyl-tropane;
- (1S,2S,3R,5R)-2-Carbomethoxy-3-(4-chlorophenyl)-tropane;
- (1S,2S,3R,5R)-2-Carbomethoxy-3-(4-methylphenyl)-tropane;
- (1S,2S,3R,5R)-2-Carbomethoxy-3-(1-naphthyl)-tropane;
- 35 (1S,2S,3R,5R)-2-Carbomethoxy-3-(4-phenylphenyl)-tropane;
- (1S,2S,3R,5R)-2-Carbomethoxy-3-(4-*t*-butyl-phenyl)-tropane;
- (1S,2S,3R,5R)-2-(4-Fluoro-benzoyl)-3-(4-fluorophenyl)-tropane
- (1S,2R,3R,5R)-2-(4-Chlorophenoxy-methyl)-3-(4-fluorophenyl)-tropane;
- (1S,2R,3R,5R)-2-(4-Chlorophenoxy-methyl)-3-(4-fluorophenyl)-tropane;

- (1S,2R,3R,5R)-2-(4-Chlorophenoxy-methyl)-3-(3,4-dichlorophenyl)-tropane;  
(1S,2R,3R,5R)-2-(4-Chlorophenoxy-methyl)-3-(4-methylphenyl)-tropane;  
(1S,2R,3R,5R)-2-(4-Benzoyloxy-methyl)-3-(4-fluorophenyl)-tropane;  
(1S,2R,3R,5R)-2-Carbomethoxy-3-(2-naphthyl)-tropane;  
5 (1S,2R,3R,5R)-2-Carbomethoxy-3-(3,4-dichlorophenyl)-tropane;  
(1S,2R,3R,5R)-2-Carbomethoxy-3-benzyl-tropane;  
(1S,2R,3R,5R)-2-Carbomethoxy-3-(4-chlorophenyl)-tropane;  
(1S,2R,3R,5R)-2-Carbomethoxy-3-(4-methylphenyl)-tropane;  
(1S,2R,3R,5R)-2-Carbomethoxy-3-(1-naphthyl)-tropane;  
10 (1S,2R,3R,5R)-2-Carbomethoxy-3-(4-phenylphenyl)-tropane;  
(1S,2R,3R,5R)-2-Carbomethoxy-3-(4-*t*-butyl-phenyl)-tropane;  
(1S,2R,3R,5R)-2-(4-Fluoro-benzoyl)-3-(4-fluorophenyl)-tropane.



**CLAIMS:**

1. A disubstituted tropane derivative of general formula (Ia) or (Ib),



or a pharmaceutically acceptable addition salt thereof or the N-oxide thereof,  
wherein

R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, or 2-hydroxyethyl;

10  $R^3$  is

- $\text{CH}_2\text{-X-R}'$ ,

wherein X is O, S, or  $\text{NR}''$ ;

wherein  $\text{R}''$  is hydrogen or alkyl; and

$\text{R}'$  is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, or  $-\text{CO-alkyl}$ ;

- 15 • **heteroaryl** which may be substituted one or more times with

- alkyl, cycloalkyl, or cycloalkylalkyl;
- phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen,  $\text{CF}_3$ , CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
- phenylphenyl;
- pyridyl which may be substituted one or more times with substituents selected from the group consisting of halogen,  $\text{CF}_3$ , CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
- thienyl which may be substituted one or more times with substituents selected from the group consisting of halogen,  $\text{CF}_3$ , CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl; or
- benzyl which may be substituted one or more times with substituents selected from the group consisting of halogen,  $\text{CF}_3$ , CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl; or

- 30 •  $(\text{CH}_2)_n\text{CO}_2\text{R}^{11}$ ,  $\text{COR}^{11}$ , or  $\text{CH}_2\text{R}^{12}$ ;

wherein  $\text{R}^{11}$  is

- alkyl, cycloalkyl, or cycloalkylalkyl;

- phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
  - phenylphenyl;
  - 5 ○ pyridyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
  - thienyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl; or
  - 10 ○ benzyl;
- n is 0 or 1; and  
R<sup>12</sup> is
- 15 ○ O-phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl; or
  - O-CO-phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
- 20 R<sup>4</sup> is
- **3,4-methylenedioxyphenyl** or
  - **phenyl, benzyl, naphthyl, or heteroaryl** all of which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, cycloalkoxy, alkyl, cycloalkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl.
- 25

2. The compound according to claim 1, wherein

R<sup>3</sup> is

- **1,2,4-oxadiazol-3-yl** which may be substituted in the 5 position with
  - 30 ○ alkyl, cycloalkyl, or cycloalkylalkyl;
  - phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
  - phenylphenyl; or
  - 35 ○ benzyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl; or
- **1,2,4-oxadiazol-5-yl** which may be substituted in the 3 position with
  - alkyl, cycloalkyl, or cycloalkylalkyl;

- phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
- phenylphenyl;
- 5 ○ benzyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl;
- pyridyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro and heteroaryl; or
- 10 ○ thienyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, alkyl, alkenyl, alkynyl, amino, nitro and heteroaryl.

15 3. The compound according to claim 1, wherein

R<sup>3</sup> is

- CH<sub>2</sub>-X-R',  
wherein X is O, S, or NR'';  
wherein R'' is hydrogen or alkyl; and
- 20 R' is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, or -CO-alkyl.

4. The compound according to any one of the claims 1-3, wherein

R<sup>4</sup> is

- phenyl, which is substituted once or twice with substituents selected from the
- 25 group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, cycloalkoxy, alkyl, cycloalkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl.

5. The compound according to claim 1, wherein

R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, or 2-hydroxyethyl;

30 R<sup>3</sup> is

- CH<sub>2</sub>-X-R',  
wherein X is O, S, or NR'';  
wherein R'' is hydrogen or alkyl; and  
R' is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, or -CO-alkyl; or
- 35 • (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sup>11</sup>, or COR<sup>11</sup>;  
wherein R<sup>11</sup> is alkyl, cycloalkyl, or cycloalkylalkyl; and n is 0 or 1;

R<sup>4</sup> is

- **phenyl** which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, cycloalkoxy, alkyl, cycloalkyl, alkenyl, alkynyl, amino, nitro, and heteroaryl.

5 6. The compound according to claim 1, wherein

R is hydrogen or alkyl;

R<sup>3</sup> is

- **CH<sub>2</sub>-X-R'**,  
wherein X is O or S; and
- **(CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sup>11</sup>**;  
wherein R<sup>11</sup> is alkyl; and n is 0 or 1;

R<sup>4</sup> is

- **phenyl** which may be substituted once or twice with substituents selected from the group consisting of halogen, CF<sub>3</sub>, and CN.

7. A compound of claim 1 which is

(1S,2R,3R,5R)-3-(3,4-Dichlorophenyl)-8-methyl-8-aza-bicyclo[3.2.1]octane-2-carboxylic acid ethyl ester;

20 (1S,2S,3R,5R)-3-(3,4-Dichlorophenyl)-8-methyl-8-aza-bicyclo[3.2.1]octane-2-carboxylic acid ethyl ester;

(1S,2R,3R,5R)-2-Hydroxymethyl-3-(3,4-dichlorophenyl)-tropane;

(1S,2S,3R,5R)-2-Hydroxymethyl-3-(3,4-dichlorophenyl)-tropane;

(1S,2R,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-tropane;

25 (1S,2S,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-tropane;

(1S,2R,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-8*H*-8-aza-bicyclo[3.2.1]octane;

(1S,2S,3R,5R)-2-Ethoxymethyl-3-(3,4-dichlorophenyl)-8*H*-8-aza-bicyclo[3.2.1]octane;

or a pharmaceutically acceptable addition salt thereof.

30 8. A pharmaceutical composition, comprising a therapeutically effective amount of a compound of any one of claims 1-7, or a pharmaceutically acceptable addition salt thereof, together with at least one pharmaceutically acceptable carrier, excipient or diluent.

35 9. The use of a compound according to any one of claims 1-7, or a pharmaceutically acceptable addition salt thereof, for the manufacture of a pharmaceutical composition for the treatment, prevention or alleviation of a disease or a disorder or a condition of a mammal, including a human, which disease, disorder or

condition is responsive to inhibition of monoamine neurotransmitter re-uptake in the central nervous system.

10.           The use according to claim 9, wherein the disease, disorder or condition is  
5 depression, pseudodementia, Ganster's syndrome, obsessive compulsive disorders, panic disorders, memory deficits, attention deficit hyperactivity disorder, obesity, anxiety and eating disorders.

11.           A method for treatment, prevention or alleviation of a disease or a disorder  
10 or a condition of a living animal body, including a human, which disorder, disease or condition is responsive to inhibition of monoamine neurotransmitter re-uptake in the central nervous system, which method comprises the step of administering to such a living animal body in need thereof a therapeutically effective amount of a compound according to any one of the claims 1-7.

15

12.           The method of claim 11, wherein the disease, disorder or condition is depression, pseudodementia, Ganster's syndrome, obsessive compulsive disorders, panic disorders, memory deficits, attention deficit hyperactivity disorder, obesity, anxiety and eating disorders.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/DK 02/00346

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07D451/02 A61K31/46

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	US 6 329 520 B1 (CARROLL F I ET AL) 11 December 2001 (2001-12-11) claims 1-5	6-12
X	WO 97 30997 A (NEUROSEARCH AS ) 28 August 1997 (1997-08-28) claims 1-9	6-12
X	WO 98 07427 A (RES TRIANGLE INST) 26 February 1998 (1998-02-26) claims 1-22	6-12
X	WO 94 04146 A (HARVARD COLLEGE) 3 March 1994 (1994-03-03) claims 26,27	6-12
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance  
"E" earlier document but published on or after the international filing date  
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
"O" document referring to an oral disclosure, use, exhibition or other means  
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art  
"&" document member of the same patent family

Date of the actual completion of the international search

26 July 2002

Date of mailing of the international search report

12.08.2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fernando Farieta

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/DK 02/00346

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 28401 A ( NEUROSEARCH AS (DK); SCHEEL KRUEG) 26 October 1995 (1995-10-26) claims 1-16	6-12
X	--- MELTZER P C ET AL: "SUBSTITUTED 3-PHENYLTROPANE ANALOGS OF COCAINE: SYNTHESIS, INHIBITION OF BINDING AT COCAINE RECOGNITION SITES, AND POSITRON EMISSION TOMOGRAPHY IMAGING" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 36, 1993, pages 855-862, XP002914108 ISSN: 0022-2623 Compound 5c	6-12
A	--- EP 0 969 005 A (LILLY CO ELI) 5 January 2000 (2000-01-05) claims 1-7	6-12
P,A	--- US 2002/004513 A1 (ANDERSSON CARL-MAGNUS A ET AL) 10 January 2002 (2002-01-10) claims 14-52	6-12
A	--- US 5 760 055 A (DAVIES HUW M L) 2 June 1998 (1998-06-02) claims 1-9	6-12
A	--- US 6 008 227 A (DAVIES HUW M L ET AL) 28 December 1999 (1999-12-28) claims 1-16	6-12
A	--- WO 99 02526 A (ORGANIX INC ) 21 January 1999 (1999-01-21) claims 1-41	6-12
A	--- WO 00 64441 A (RESPIRATORIUS AB ) 2 November 2000 (2000-11-02) claims 1-17 -----	6-12

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/DK 02/00346

### Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **11 and 12**  
because they relate to subject matter not required to be searched by this Authority, namely:  
**see FURTHER INFORMATION sheet PCT/ISA/210**
2. ☒ Claims Nos.: **1-5**  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
**see FURTHER INFORMATION sheet PCT/ISA/210**
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.



**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

**Continuation of Box I.1**

**Claims Nos.: 11 and 12**

Claims 11-12 relate to methods of treatment of the human or animal body by surgery or by therapy/ diagnostic methods practised on the human or animal body/ Rule 39.1.(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.

-----

**Continuation of Box I.2**

**Claims Nos.: 1-5**

Present claims 1-5 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts related to the compounds of claim 6-7.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 02/00346

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 6329520	B1	11-12-2001	
		US 5736123 A	07-04-1998
		US 5413779 A	09-05-1995
		US 5496953 A	05-03-1996
		US 5128118 A	07-07-1992
		AU 4075999 A	13-12-1999
		EP 1079833 A1	07-03-2001
		JP 2002516278 T	04-06-2002
		WO 9961023 A1	02-12-1999
		AU 4232797 A	06-03-1998
		EP 0993301 A1	19-04-2000
		JP 2001525795 T	11-12-2001
		WO 9807427 A1	26-02-1998
		US 5935953 A	10-08-1999
		US 6123917 A	26-09-2000
		AT 182794 T	15-08-1999
		AU 668371 B2	02-05-1996
		AU 3064592 A	15-06-1993
		CA 2123570 A1	27-05-1993
		DE 69229744 D1	09-09-1999
		DE 69229744 T2	27-04-2000
		DK 644775 T3	06-03-2000
		EP 0644775 A1	29-03-1995
		EP 0897922 A2	24-02-1999
		EP 0905135 A2	31-03-1999
		ES 2138601 T3	16-01-2000
		FI 942231 A	13-05-1994
		GR 3031579 T3	31-01-2000
		JP 2852467 B2	03-02-1999
		JP 7501074 T	02-02-1995
		KR 251339 B1	01-09-2000
		NO 941753 A	10-05-1994
		WO 9309814 A1	27-05-1993
		US 5380848 A	10-01-1995
		AU 650059 B2	09-06-1994
		AU 8519591 A	02-03-1992
		DE 69123151 D1	19-12-1996
		EP 0542903 A1	26-05-1993
		GR 3022529 T3	31-05-1997
		JP 6503556 T	21-04-1994
		AT 145140 T	15-11-1996
		CA 2089070 A1	10-02-1992
		DK 542903 T3	21-04-1997
		ES 2099167 T3	16-05-1997
		WO 9202260 A1	20-02-1992
WO 9730997	A	28-08-1997	
		AT 203023 T	15-07-2001
		AU 720358 B2	01-06-2000
		AU 1794097 A	10-09-1997
		BG 102637 A	30-06-1999
		BR 9707636 A	27-07-1999
		CA 2244773 A1	28-08-1997
		CN 1211982 A ,B	24-03-1999
		CZ 9802520 A3	11-11-1998
		DE 69705608 D1	16-08-2001
		DE 69705608 T2	16-05-2002
		DK 885220 T3	15-10-2001
		EE 9800254 A	15-02-1999
		WO 9730997 A1	28-08-1997

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 02/00346

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9730997	A		EP 1130020 A1	05-09-2001
			EP 0885220 A1	23-12-1998
			HU 9901199 A2	30-08-1999
			JP 3238414 B2	17-12-2001
			JP 2000504739 T	18-04-2000
			NO 983877 A	21-08-1998
			NZ 330886 A	25-02-1999
			PL 328503 A1	01-02-1999
			PT 885220 T	30-11-2001
			RU 2167876 C2	27-05-2001
			SI 885220 T1	31-12-2001
			SK 92998 A3	04-11-1998
			TR 9801641 T2	23-11-1998
			US 6288079 B1	11-09-2001
			US 2001018444 A1	30-08-2001
			ZA 9701525 A	21-10-1997
WO 9807427	A	26-02-1998	US 5935953 A	10-08-1999
			AU 4232797 A	06-03-1998
			EP 0993301 A1	19-04-2000
			JP 2001525795 T	11-12-2001
			WO 9807427 A1	26-02-1998
			US 6329520 B1	11-12-2001
WO 9404146	A	03-03-1994	AU 5088993 A	15-03-1994
			WO 9404146 A1	03-03-1994
			US 5506359 A	09-04-1996
			US 5770180 A	23-06-1998
WO 9528401	A	26-10-1995	AU 690257 B2	23-04-1998
			AU 2257595 A	10-11-1995
			BG 63259 B1	31-07-2001
			BG 100883 A	31-03-1998
			BR 9507489 A	12-08-1997
			CA 2187309 A1	26-10-1995
			CN 1148854 A ,B	30-04-1997
			CZ 9602982 A3	11-06-1997
			EE 9600132 A	15-04-1997
			WO 9528401 A1	26-10-1995
			EP 0756596 A1	05-02-1997
			FI 965074 A	17-12-1996
			HU 75865 A2	28-05-1997
			JP 2899418 B2	02-06-1999
			JP 9505607 T	03-06-1997
			KR 210417 B1	15-07-1999
			LV 11738 A	20-04-1997
			LV 11738 B	20-12-1997
			NO 964180 A	16-12-1996
			NZ 284075 A	26-01-1998
			PL 316876 A1	17-02-1997
			RU 2134264 C1	10-08-1999
			SK 128896 A3	09-04-1997
			US 5736556 A	07-04-1998
			ZA 9508760 A	09-05-1996
EP 0969005	A	05-01-2000	AU 4819099 A	05-01-2000
			EP 0969005 A1	05-01-2000
			JP 2002518331 T	25-06-2002

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 02/00346

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0969005	A	WO 9965492 A1 US 6107307 A	23-12-1999 22-08-2000
US 2002004513	A1	10-01-2002 AU 4007201 A WO 0166521 A1	17-09-2001 13-09-2001
US 5760055	A	02-06-1998 US 6008227 A US 5262428 A US 5763455 A AU 673265 B2 AU 6767194 A CA 2163095 A1 EP 0697871 A1 JP 2906085 B2 JP 8510238 T WO 9426274 A1 AU 668721 B2 AU 3932893 A CA 2131789 A1 EP 0630375 A1 JP 7504665 T JP 2578727 B2 WO 9318033 A1 US 5342949 A	28-12-1999 16-11-1993 09-06-1998 31-10-1996 12-12-1994 24-11-1994 28-02-1996 14-06-1999 29-10-1996 24-11-1994 16-05-1996 05-10-1993 16-09-1993 28-12-1994 25-05-1995 05-02-1997 16-09-1993 30-08-1994
US 6008227	A	28-12-1999 US 5262428 A AU 673265 B2 AU 6767194 A CA 2163095 A1 EP 0697871 A1 JP 2906085 B2 JP 8510238 T WO 9426274 A1 US 5760055 A US 5763455 A AU 668721 B2 AU 3932893 A CA 2131789 A1 EP 0630375 A1 JP 7504665 T JP 2578727 B2 WO 9318033 A1 US 5342949 A	16-11-1993 31-10-1996 12-12-1994 24-11-1994 28-02-1996 14-06-1999 29-10-1996 24-11-1994 02-06-1998 09-06-1998 16-05-1996 05-10-1993 16-09-1993 28-12-1994 25-05-1995 05-02-1997 16-09-1993 30-08-1994
WO 9902526	A	21-01-1999 US 5948933 A EP 0996619 A1 JP 2001509508 T WO 9902526 A1 US 6353105 B1 US 6417221 B1 US 6291512 B1	07-09-1999 03-05-2000 24-07-2001 21-01-1999 05-03-2002 09-07-2002 18-09-2001
WO 0064441	A	02-11-2000 AU 5259100 A EP 1173168 A2 WO 0064441 A2 AU 5861900 A CN 1355698 T EP 1185263 A2	10-11-2000 23-01-2002 02-11-2000 02-01-2001 26-06-2002 13-03-2002

## INTERNATIONAL SEARCH REPORT

### Information on patent family members

**International Application No**

**PCT/DK 02/00346**

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0064441	A	WO 0076500 A2	21-12-2000
-----			